arXiv:1212.2335v1 [math.AP] 11 Dec 2012

SOME NEW REGULARITY CRITERIA FOR THE 3D NAVIER-STOKES EQUATIONS

DAOYUAN FANG CHENYIN QIAN
DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY,
HANGZHOU, 310027, CHINA

Abstract. Several types of new regularity criteria for Leray-Hopf weak solutions u to the 3D Navier-Stokes equations are obtained. Some of them are based on the third component u_3 of velocity under Prodi-Serrin index condition, another type is in terms of ω_3 and $\partial_3 u_3$ with Prodi-Serrin index condition. And a very recent work of the authors, based on only one of the nine entries of the gradient tensor, is renovated.

1. Introduction

In the present paper, we address sufficient conditions for the regularity of weak solutions of the Cauchy problem for the Navier-Stokes equations in $\mathbb{R}^3 \times (0,T)$:

$$\begin{cases} \dfrac{\partial u}{\partial t} - \nu \Delta u + (u\cdot\nabla)u + \nabla p = 0, & \text{in } \mathbb{R}^3 \times (0,T), \\ \nabla \cdot u = 0, & \text{in } \mathbb{R}^3 \times (0,T), \\ u(x,0) = u_0, \quad \text{in } \mathbb{R}^3, & \end{cases} \tag{1.1}$$

where $u = (u_1, u_2, u_3) : \mathbb{R}^3 \times (0,T) \to \mathbb{R}^3$ is the velocity field, $p : \mathbb{R}^3 \times (0,T) \to \mathbb{R}^3$ is a scalar pressure, and u_0 is the initial velocity field, $\nu > 0$ is the viscosity. We set $\nabla_h = (\partial_{x_1}, \partial_{x_2})$ as the horizontal gradient operator and $\Delta_h = \partial_{x_1}^2 + \partial_{x_2}^2$ as the horizontal Laplacian, and Δ and ∇ are the usual Laplacian and the gradient operators, respectively. Here we use the classical notations

$$(u\cdot\nabla)v = \sum_{i=1}^{3} u_i \partial_{x_i} v_k, \ (k = 1,2,3), \quad \nabla \cdot u = \sum_{i=1}^{3} \partial_{x_i} u_i,$$

and for sake of simplicity, we denote ∂_{x_i} by ∂_i.

It is well known that the weak solution of the Navier-Stokes equations (1.1) is unique and regular in two dimensions. However, in three dimensions, the regularity problem of weak solutions of Navier-Stokes equations is an outstanding open problem in mathematical fluid mechanics. The weak solutions are known to exist globally in time, but the uniqueness, regularity, and continuous dependence on initial data for weak solutions are still open problems. Furthermore, strong solutions in the 3D case are known to exist for a short interval of time whose length depends on the initial data. Moreover, this strong solution is known to be unique and to depend continuously on the initial data (see, for example, [22], [24]). Let us recall the definition of Leray-Hopf weak solution. We set

$$\mathcal{V} = \{\phi : \ \text{the 3D vector valued } C_0^\infty \text{ functions and } \nabla\cdot\phi = 0\},$$

2000 Mathematics Subject Classification. 35Q30; 76D05.
Key words. 3D Navier-Stokes equations; Leray-Hopf weak solution; Regularity criterion.
*E-mail addresses: dyf@zju.edu.cn (D. Fang), qcyjcsx@163.com(C. Qian).

which will form the space of test functions. Let H and V be the closure spaces of $\mathcal{V}$ in L^2 under L^2-topology, and in H^1 under H^1-topology, respectively.

For $u_0 \in H$, the existence of weak solutions of (1.1) was established by Leray [15] and Hopf in [11], that is, u satisfies the following properties:
(i) $u \in C_w([0,T);H) \cap L^2(0,T;V)$, and $\partial_t u \in L^1(0,T;V')$, where V' is the dual space of V;
(ii) u verifies (1.1) in the sense of distribution, i.e., for every test function $\phi \in C_0^\infty([0,T);\mathcal{V})$, and for almost every $t, t_0 \in (0,T)$, we have

$$\begin{aligned}\int_{\mathbb{R}^3} u(x,t)\cdot\phi(x,t)dx &- \int_{\mathbb{R}^3} u(x,t_0)\cdot\phi(x,t_0)dx\\ &= \int_{t_0}^t \int_{\mathbb{R}_3} [u(x,t)\cdot(\phi_t(x,t)+\nu\Delta\phi(x,t))]dxds\\ &\quad + \int_{t_0}^t \int_{\mathbb{R}_3} [(u(x,t)\cdot\nabla)\phi(x,t)]\cdot u(x,t))]dxds\end{aligned}$$

(iii) The energy inequality, i.e.,

$$\|u(\cdot,t)\|_{L^2}^2 + 2\nu \int_{t_0}^t \|\nabla u(\cdot,s)\|_{L^2}^2 ds \le \|u_0\|_{L^2}^2,$$

for every t and almost every t_0.

It is well known, if $u_0 \in V$, a weak solution becomes strong solution of (1.1) on $(0,T)$ if, in addition, it satisfies

$$u \in C([0,T);V) \cap L^2(0,T;H^2) \text{ and } \partial_t u \in L^2(0,T;H).$$

We know the strong solution is regular(say, classical) and unique (see, for example, [22], [24]).

Researchers are interested in the classical problem of finding sufficient conditions for weak solutions of (1.1) such that the weak solutions become regular, and the first result is usually referred as Prodi-Serrin conditions (see [20] and [21]), which states that if a weak solution u is in the class of

$$u \in L^t(0,T;L^s(\mathbb{R}^3)), \quad \frac{2}{t}+\frac{3}{s}=1, \ s\in[3,\infty], \tag{1.2}$$

then the weak solution becomes regular. Recently, H. Bae and H. Choe in [1] gave a two components Prodi-Serrin index criterion. Up to now, there are many results show that one can use only one component (say u_3) to determine the regularity of u. Say, I. Kukavica and M. Ziane in [12] proved a regularity criterion under the following condition

$$u_3 \in L^t(0,T;L^s(\mathbb{R}^3)), \quad \frac{2}{t}+\frac{3}{s}\le\frac{5}{8}, \ s\in[\frac{24}{5},\infty].$$

Then, it was improved by C. Cao and E.Titi in [6] to

$$u_3 \in L^t(0,T;L^s(\mathbb{R}^3)), \quad \frac{2}{t}+\frac{3}{s}\le\frac{2(s+1)}{3s}, \ s\in(\frac{7}{2},\infty].$$

And then, Y. Zhou and M. Pokorný in [26] changed the regular criterion to

$$u_3 \in L^t(0,T;L^s(\mathbb{R}^3)), \quad \frac{2}{t}+\frac{3}{s}\le\frac{3}{4}+\frac{1}{2s}, \ s\in(\frac{10}{3},\infty].$$

More relative results, we refer to [16], [25] and the reference there in. One can see that the above mentioned results on u_3 cannot satisfy the Prodi-Serrin index condition, and it seems to be a price when one reduce the components of u to one. It is nature to think about what supplement is necessary to insure the Prodi-Serrin condition based on one velocity component. For example, P. Penel and M. Pokorný in [18] proved the u was regular, if

$$u_3 \in L^\beta(0,T;L^\alpha(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha}+\frac{2}{\beta} \leq 1, \alpha \in (3,\infty], \beta \in [2,\infty),$$

and one of the following conditions holds true:
(a) $\partial_3 u_1, \partial_3 u_2$ belong to $L^\beta(0,T;L^\alpha(\mathbb{R}^3))$ with $\frac{3}{\alpha}+\frac{2}{\beta} \leq 2, \alpha \in (3/2,\infty], \beta \in [1,\infty)$;
(b) $\partial_2 u_1, \partial_1 u_2$ belong to $L^\beta(0,T;L^\alpha(\mathbb{R}^3))$ with $\frac{3}{\alpha}+\frac{2}{\beta} \leq 2, \alpha \in [2,3], \beta \in [2,\infty]$;
(c) $\partial_3 u_2$ belong to $L^\beta(0,T;L^\alpha(\mathbb{R}^3))$ with $\frac{3}{\alpha}+\frac{2}{\beta} \leq 2, \alpha \in (3/2,\infty], \beta \in [1,\infty)$, and $\partial_2 u_1$ belong to $L^\beta(0,T;L^\alpha(\mathbb{R}^3))$ with $\frac{3}{\alpha}+\frac{2}{\beta} \leq 2, \alpha \in [2,3], \beta \in [2,\infty]$.
Moreover, the authors also mentioned in the Remark 2 in [18], the condition in (a) can be replaced by $\partial_3 u_2,\ \partial_2 u_2$, or $\partial_3 u_2,\ \partial_1 u_1$, or $\partial_3 u_1,\ \partial_2 u_2$, or $\partial_3 u_1,\ \partial_2 u_1$. Similarly, in (c) one can replace $\partial_3 u_2$ by $\partial_3 u_1$, and replace $\partial_2 u_1$ by $\partial_1 u_2$ respectively.

From the above , we can see that the assumptions on derivative component did not contain $\partial_i u_3$ $i=1,2,3$. One purpose of this paper is to make up this work, by using the incompressibility condition, we give an estimate on velocity, which is different from [18] and then get a regularity criterion on $\partial_i u_3$ $i=1,2,3$, for detail see the proof of Theorem 1.3 below. On the other hand, similar to (a), (b) or (c), we also consider cases of the given conditions in terms of only one component $\partial_i u_j$ of ∇u such that u_3 satisfies the Prodi-Serrin condition in Theorem 1.4 and Corollary 1.6.

H. Beirão da Veiga in [2] obtained a sufficient condition for regularity using the vorticity rather than the velocity. His result says that if the vorticity $\omega = curl u$ satisfies

$$\omega \in L^t(0,T;L^s(\mathbb{R}^3)), \quad \frac{2}{t}+\frac{3}{s} \leq 2, \ s \in (1,\infty),$$

then u becomes the strong solution. Later, D. Chae, H. Choe proved that it was sufficient to control only two components of the vorticity vector in [7]. Another goal of this paper is to prove the regularity of u by imposing the Prodi-Serrin condition on only one component of vorticity (say ω_3). Precisely, we get a regularity criterion with Prodi-Serrin condition, by using only ω_3 and $\partial_3 u_3$, which is shown in Theorem 1.1.

Besides, we would like to point out that the full regularity of weak solutions can also be proved under alternative assumptions on the gradient of the velocity ∇u, for instance

$$\nabla u \in L^t(0,T;L^s(\mathbb{R}^3)), \quad \frac{2}{t}+\frac{3}{s} = 2, \ s \in [\frac{3}{2},\infty]. \tag{1.3}$$

Enlightened by above, we also want to get some better regularity criteria which are also coincident with the standard Prodi-Serrin condition based on some components of ∇u. To begin with, we mention some results in this direction at first, P. Penel and M. Pokorný in [18] proved that if

$$\partial_3 u \in L^\beta(0,T;L^\alpha(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha}+\frac{2}{\beta} \leq \frac{3}{2}, \alpha \in [2,\infty], \beta \in [1,\infty).$$

then the weak solution was regular. After that many authors improved this result, such as I. Kukavica and M. Ziane in [13] considered the case of the condition

$$\partial_3 u \in L^\beta(0,T;L^\alpha(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha}+\frac{2}{\beta} \leq 2, \alpha \in [\frac{9}{4},3].$$

As to the gradient of one velocity component ∇u_3, M. Pokorný in [19] proved the weak solution was actually regular if ∇u_3 satisfied

$$\nabla u_3 \in L^\beta(0,T;L^\alpha(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha}+\frac{2}{\beta}\leq\frac{3}{2}, \alpha\in[2,\infty].$$

Y. Zhou and M. Pokorný in [26] improved the result to

$$\nabla u_3 \in L^\beta(0,T;L^\alpha(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha}+\frac{2}{\beta}\leq\begin{cases}\frac{19}{12}+\frac{1}{2\alpha}, \alpha\in(\frac{30}{19},3]\\ \frac{3}{2}+\frac{3}{4\alpha}, \alpha\in(3,\infty],\end{cases}$$

moreover, Y. Zhou and M. Pokorný also proved a improved result, for more detail we refer to [27]. Motivated by the above, we consider the case of two gradient velocity components and one of them satisfies the Prodi-Serrin condition, see Theorem 1.7 and Corollary 1.8. We shall point out that two gradient velocity components are not all the diagonal elements, this is more difficult than the diagonal case, for the detail see Remark 1.9 below.

Recently, the regularity criterion in terms of only one of the gradient tensor was gotten by C.S. Cao, E.S. Titi in [5] under the assumptions

$$\frac{\partial u_j}{\partial x_k}\in L^\beta(0,T;L^\alpha(\mathbb{R}^3)), \text{ when } j\neq k$$

$$\text{and where } \alpha>3, 1\leq\beta<\infty, \text{and } \frac{3}{\alpha}+\frac{2}{\beta}\leq\frac{\alpha+3}{2\alpha}, \tag{1.4}$$

or

$$\frac{\partial u_j}{\partial x_j}\in L^\beta(0,T;L^\alpha(\mathbb{R}^3)),$$

$$\text{and where } \alpha>2, 1\leq\beta<\infty, \text{and } \frac{3}{\alpha}+\frac{2}{\beta}\leq\frac{3(\alpha+2)}{4\alpha}. \tag{1.5}$$

In [9], we improved this result. And here, we again study it and get an improvement of the results of [9], which is shown in Theorem 1.10. However, it is also noted that the above conditions are not coincident with the Prodi-Serrin condition.

Now, we list our main results as follows:

Theorem 1.1. *Let u be a Leray-Hopf weak solution to the 3D Navier-Stokes equations* (1.1) *with the initial value $u_0\in V$. Suppose*

$$\omega_3, \partial_3 u_3\in L^\beta(0,T;L^\alpha(\mathbb{R}^3)) \tag{1.6}$$

with

$$\frac{3}{\alpha}+\frac{2}{\beta}\leq 2, \alpha\in(\frac{3}{2},\infty], \beta\in[2,\infty), \tag{1.7}$$

where $\omega_3=\partial_1u_2-\partial_2u_1$ is the third component of the vorticity $\omega=curl u$.
Then u is regular.

Remark 1.2. As mentioned above, the authors in [18] proved the regularity criterion with u_3, ∂_1u_2, and ∂_2u_1. Can we replace $\partial_1u_2, \partial_2u_1$ by w_3? It is open.

Theorem 1.3. *Let u_0 and u be as in Theorem 1.1. Suppose*

$$u_3 \in L^{\beta}(0,T;L^{\alpha}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_1}+\frac{2}{\beta_1}\le 1, \alpha_1\in(3,\infty], \beta_1\in[2,\infty), \tag{1.8}$$

and one of the following conditions holds:

$$(i)\ \partial_3 u_2, \partial_3 u_3 \in L^{\beta_2}(0,T;L^{\alpha_2}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_2}+\frac{2}{\beta_2}\le 2, \alpha_2\in[2,3], \beta_2\in[2,\infty); \tag{1.9}$$

$$(ii)\ \partial_3 u_1, \partial_3 u_3 \in L^{\beta_2}(0,T;L^{\alpha_2}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_2}+\frac{2}{\beta_2}\le 2, \alpha_2\in[2,3], \beta_2\in[2,\infty). \tag{1.10}$$

Then u is regular.

Theorem 1.4. *Let u be a Leray-Hopf weak solution to the 3D Navier-Stokes equations* (1.1) *with the initial value $u_0 \in V$. For any i, j with $1 \le i \le 3$ and $1 \le j \le 2$, suppose*
(a) $i \neq j$,

$$u_3 \in L^{\beta_1}(0,T;L^{\alpha_1}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_1}+\frac{2}{\beta_1}\le 1, \alpha_1\in(3,\infty], \beta_1\in[2,\infty), \tag{1.11}$$

and

$$\partial_i u_j \in L^{\beta_2}(0,T;L^{\alpha_2}(\mathbb{R}^3)), \tag{1.12}$$

with

$$\frac{3}{\alpha_2}+\frac{2}{\beta_2}\le \frac{\alpha_2+1}{\alpha_2},\ \ \alpha_2\in(2,\infty), \beta_2\in(1,\infty). \tag{1.13}$$

(b) $i = j$, u_3 satisfies the condition (1.11), *and*

$$\partial_j u_j \in L^{\beta_3}(0,T;L^{\alpha_3}(\mathbb{R}^3)), \tag{1.14}$$

with

$$\frac{3}{2\alpha_3}+\frac{2}{\beta_3}\le f(\alpha_3),\ \ \alpha_3\in[2,\infty), \beta_3\in(1,\infty) \tag{1.15}$$

where

$$f(\alpha_3)=\frac{\sqrt{24\alpha_3^2-24\alpha_3+9}-2\alpha_3}{2\alpha_3}.$$

Then u is regular.

Remark 1.5. Recall that in [9], we proved the regularity criterion only on one component $\partial_i u_j, i, j = 1, 2, 3$. We note that the condition is more restrictive than (1.13) and (1.15). This means that the condition on $\partial_i u_j, i, j = 1, 2, 3$. becomes weaker with the help of the condition (1.11).

Corollary 1.6. *Suppose that $u_0 \in V$, and u is a Leray-Hopf weak solution to the 3D Navier-Stokes equations* (1.1). *For any i, j with $1 \le i \le 3$ and $1 \le j \le 2$, suppose*
(a) $i \neq j$, u_3 satisfies the condition (1.11) *and*

$$\partial_i u_j \in L^{\beta_2}(0,T;L^{\alpha_2}(\mathbb{R}^3)),\ \frac{3}{\alpha_2}+\frac{2}{\beta_2}\le 1,\ \ \alpha_2\in(2,\infty), \beta_2\in(1,\infty) \tag{1.16}$$

(b) $i = j$, u_3 satisfies the condition (1.11), *and*

$$\partial_j u_j \in L^{\beta_3}(0,T;L^{\alpha_3}(\mathbb{R}^3)), \frac{3}{\alpha_3}+\frac{2}{\beta_3}\le \frac{3}{2},\ \ \alpha_3\in[2,6], \beta_3\in[2,\infty). \tag{1.17}$$

Then u is regular.

If we substitute the condition on u_3 by the component of the gradient of the velocity, we have the following regularity criterion, which is a further improvement of the above mentioned results of [18].

Theorem 1.7. *Let u be a Leray-Hopf weak solution to the 3D Navier-Stokes equations* (1.1) *with the initial value $u_0 \in V$. Suppose*

$$\partial_3 u_i \in L^{\beta_1}(0,T;L^{\alpha_1}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_1}+\frac{2}{\beta_1} \le 2, \alpha_1 \in [2,3], \beta_1 \in [2,\infty), i=1 \text{ or } 2, \quad (1.18)$$

and

$$\partial_3 u_3 \in L^{\beta_2}(0,T;L^{\alpha_2}(\mathbb{R}^3)), \quad (1.19)$$

with

$$\frac{3}{2\alpha_2}+\frac{2}{\beta_2} \le f(\alpha_2),\ \ \alpha_2 \in [2,3], \beta_2 \in [1,\infty) \quad (1.20)$$

where

$$f(\alpha_2)=\frac{\sqrt{24\alpha_2^2-24\alpha_2+9}-2\alpha_2}{2\alpha_2}.$$

Then u is regular.

Corollary 1.8. *Suppose that $u_0 \in V$, and u is a Leray-Hopf weak solution to the 3D Navier-Stokes equations* (1.1)*. Assume*

$$\partial_3 u_i \in L^{\beta_1}(0,T;L^{\alpha_1}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_1}+\frac{2}{\beta_1} \le 2, \alpha_1 \in [2,3], \beta_1 \in [2,\infty), i=1 \text{ or } 2, \quad (1.21)$$

and

$$\partial_3 u_3 \in L^{\beta_2}(0,T;L^{\alpha_2}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_2}+\frac{2}{\beta_2} \le \frac{3}{2}, \alpha_2 \in [2,3], \beta_2 \in [4,\infty). \quad (1.22)$$

Then u is regular.

Remark 1.9. Here we only need two components of the gradient of the velocity and one of them is not on the diagonal elements of ∇u. On the case of the diagonal elements of ∇u, P. Penel and M. Pokorný in [18] proved the u is regular when

$$\partial_2 u_2, \partial_3 u_3 \in L^{\beta}(0,T;L^{\alpha}(\mathbb{R}^3)),$$

and α, β satisfied

$$\frac{3}{\alpha}+\frac{2}{\beta} \le 2, \alpha \in (\frac{3}{2},\infty], \beta \in [2,\infty).$$

Moreover, the condition on $\partial_3 u_i$ satisfies the Prodi-Serrin condition, which is an improvement of the result of P. Penel and M. Pokorný in [19]. Finally, we note that $\partial_3 u_i$, $i=1$ or 2, is not the diagonal element of ∇u. Thus, we cannot use the method of by multiplying u_i to the ith equation of (1.1) to get the form $\partial_3 u_i$, $i=1$ or 2. Therefore, it is more difficult to get the regularity criterion based on $\partial_i u_i$ and $\partial_j u_j$, $i,j \in \{1,2,3\}$ with $i \neq j$.

Theorem 1.10. *Let u_0 and u be as in Theorem 1.1. Suppose in addition*

$$\partial_k u_k \in L^\beta(0,T;L^\alpha(\mathbb{R}^3)),\ \textit{for some } k\in\{1,2,3\} \tag{1.23}$$

with

$$\frac{3}{2\alpha}+\frac{2}{\beta}\le g(\alpha),\ \alpha\in(\frac{9}{5},\infty),\ \beta\in(1,\infty), \tag{1.24}$$

where

$$g(\alpha)=\frac{\sqrt{289\alpha^2-264\alpha+144}-7\alpha}{8\alpha}.$$

Then u is regular.

Remark 1.11. This theorem is an improvement of [5] and [26] (see figure 1 below), and is also an improvement of theorem 1.2 (i) and theorem 1.3 in [9]. Moreover, we point out that the first part of Theorem 1.2 in [9], can be simplified to the following form:
For $j\neq k$, suppose that u satisfies

$$\int_0^T \|\partial_j u_k\|_\alpha^\beta d\tau\le M,\ \text{for some } M>0, \tag{1.25}$$

with

$$\frac{3}{2\alpha}+\frac{2}{\beta}\le f(\alpha),\ \alpha\in(3,\infty)\text{ and } 1\le\beta<\infty, \tag{1.26}$$

where

$$f(\alpha)=\frac{\sqrt{103\alpha^2-12\alpha+9}-9\alpha}{2\alpha}, \tag{1.27}$$

then u is regular. This statement shows the same line as in the Figure 1. in [9]. In the proof of Theorem 1.2 (i) of [9], if we substitute σ_1 and β by

$$\frac{1}{\sigma_1}=\frac{(7-\frac{3}{\alpha})+\sqrt{\frac{9}{\alpha^2}-\frac{12}{\alpha}+103}}{18},\ \beta=\frac{2\sigma_1}{9-8\sigma_1},$$

we can get the desired result.

For convenience, we recall the following version of the three-dimensional Sobolev and Ladyzhenskaya inequalities in the whole space $\mathbb{R}^3$ (see, for example, [8], [10], [14]). There exists a positive constant C such that

$$\begin{aligned}\|u\|_r &\le C\|u\|_2^{\frac{6-r}{2r}}\|\partial_1 u\|_2^{\frac{r-2}{2r}}\|\partial_2 u\|_2^{\frac{r-2}{2r}}\|\partial_3 u\|_2^{\frac{r-2}{2r}}\\ &\le C\|u\|_2^{\frac{6-r}{2r}}\|\nabla u\|_2^{\frac{3(r-2)}{2r}},\end{aligned} \tag{1.28}$$

for every $u\in H^1(\mathbb{R}^3)$ and every $r\in[2,6]$, where C is a constant depending only on r. Taking ∇div on both sides of (1.1) for smooth $(u;p)$, one can obtain

$$-\Delta(\nabla p)=\sum_{i,j}^{3}\partial_i\partial_j(\nabla(u_iu_j)),$$

therefore, the Calderon-Zygmund inequality in $\mathbb{R}^3$ (see [23])

$$\|\nabla p\|_q\le C\||\nabla u||u|\|_q, 1<q<\infty, \tag{1.29}$$

holds, where C is a positive constant depending only on q. And there is another estimates for the pressure

$$\|p\|_q\le C\|u\|_{2q}^2,\ 1<q<\infty, \tag{1.30}$$



FIGURE 1. Case of $j = k$

The line "**(1)**" is the result of C.S. Cao, E.S. Titi in [5]. The line "**(3)**" is our result, which mean (1.24). The result of Y. Zhou, M. Pokorný in [26] is showed by line "**(2)**".

Recall also that if div$u = 0$ then the vorticity $\omega =$curl$u= \nabla\times u$ has the following estimates (see [18]):

$$C\|\omega\|_q \leq \|\nabla u\|_q \leq C(q)\|\omega\|_q,\ 1 < q < \infty. \tag{1.31}$$

Moreover, if div$u = 0$, the expression

$$\Delta u = \nabla(\mathrm{div} u) - \nabla \times (\nabla \times u) \tag{1.32}$$

can be reduced to

$$\Delta u = -\nabla \times (\nabla \times u) = -\nabla \times \omega. \tag{1.33}$$

On the other hand, note that div$\omega = 0$, applying (1.31), we have

$$C\|\nabla \times \omega\|_q \leq \|\nabla\omega\|_q \leq C(q)\|\nabla \times \omega\|_q,\ 1 < q < \infty. \tag{1.34}$$

Therefore, by (1.31) and (1.33), we have

$$C\|\Delta u\|_q \leq \|\nabla\omega\|_q \leq C(q)\|\Delta u\|_q,\ 1 < q < \infty. \tag{1.35}$$

If u be a divergence-free sufficiently smooth vector field in $\mathbb{R}^3$, there exist a C such that (see [17] Lemma 2.1)

$$\|\partial_i u_j\|_q \leq C(\|\omega_3\|_q + \|\partial_3 u_3\|_q),\ 1 < q < \infty,\ i, j = 1, 2. \tag{1.36}$$

2. PROOF OF MAIN RESULTS

In this section, under the assumptions of Theorems 1.1-1.4, Theorem 1.7 or Theorem1.10 in Section 1 respectively, we prove our main results. First of all, we note that, by the energy inequality, for Leray-Hopf weak solutions, we have (see, for example, [22], [24] for detail)

$$\|u(\cdot, t)\|_{L^2}^2 + 2\nu \int_0^t \|\nabla u(\cdot, s)\|_{L^2}^2 ds \leq K_1, \tag{2.1}$$

for all $0 < t < T$, where $K_1 = \|u_0\|_{L^2}^2$.

It is well known that there exists a unique strong solution u local in time if $u_0 \in V$. In addition, this strong solution $u \in C([0, T^*); V) \cap L^2(0, T^*; H^2(\mathbb{R}^3))$ is the only weak solution with the initial datum u_0, where $(0, T^*)$ is the maximal interval of existence of the unique strong solution. If $T^* \geq T$, then there is nothing to prove. If, on the other hand, $T^* < T$, then our strategy is to show that the H^1 norm of this strong solution is bounded uniformly in time over the interval $(0, T^*)$, provided additional conditions in Theorem 1.1-1.4, Theorem 1.7 or Theorem1.10 in Section 1 are valid. As a result the interval $(0, T^*)$ can not be a maximal interval of existence, and consequently $T^* \geq T$, which concludes our proof.

In order to prove the H^1 norm of the strong solution u is bounded on interval $(0, T^*)$, combing with the energy equality (2.1), it is sufficient to prove

$$\|\nabla u\|_{L^2}^2 + \nu \int_0^t \|\Delta u\|_{L^2}^2 d\tau \leq C, \ \forall\ t \in (0, T^*) \tag{2.2}$$

where the constant C depends on T, K_1. We recall the following lemma (see [4]), which is useful for our proof of the Theorems.

Lemma 2.1. *Assume $v \in L_t^\infty L_x^2(\mathbb{R}^3 \times I)$ and $\nabla v \in L_t^2 L_x^2(\mathbb{R}^3 \times I)$, where I is an open interval. Then $v \in L_t^s L_x^r(\mathbb{R}^3 \times I)$ for all r and s such that*

$$\frac{2}{s} + \frac{3}{r} = \frac{3}{2} \text{ with } 2 \leq r \leq 6.$$

and it holds

$$\|v\|_{L_t^s L_x^r} \leq C \|v\|_{L_t^\infty L_x^2}^{\frac{6-r}{2r}} \|\nabla v\|_{L_t^2 L_x^2}^{\frac{3(r-2)}{2r}}.$$

Proof of Theorem 1.1 It is sufficient to prove the case of

$$\omega_3, \partial_3 u_3 \in L^\beta(0, T; L^\alpha(\mathbb{R}^3))$$

with

$$\frac{3}{\alpha} + \frac{2}{\beta} = 2, \alpha \in (\frac{3}{2}, \infty], \beta \in [2, \infty),$$

and we point out that in the following Theorems, we only need to prove the "=" case. Taking the curl on (1.1), we obtain

$$\frac{\partial \omega}{\partial t} - \nu \Delta \omega + (u \cdot \nabla)\omega - (\omega \cdot \nabla) u = 0.$$

We taking the inner product of above inequality with ω in $L^2(\mathbb{R}^3)$, the vorticity ω satisfies

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|\omega\|_{L^2}^2 + \nu \|\nabla \omega\|_{L^2}^2 &= \int_{\mathbb{R}^3} (\omega \cdot \nabla) u \cdot \omega dx = \sum_{i,j=1}^3 \int_{\mathbb{R}^3} \omega_i \partial_i u_j \omega_j dx \\ &= \sum_{i,j=1}^2 \int_{\mathbb{R}^3} \omega_i \partial_i u_j \omega_j dx + \sum_{i=1}^2 \int_{\mathbb{R}^3} \omega_i \partial_i u_3 \omega_3 dx \\ &\quad + \sum_{i=1}^3 \int_{\mathbb{R}^3} \omega_3 \partial_3 u_i \omega_i dx \\ &= I_1(t) + I_2(t) + I_3(t). \end{aligned} \tag{2.3}$$

We estimate $I_i(t),\ i=1,2,3$ in turn. For the first term, apply Hölder's inequality, we obtain

$$\begin{aligned} |I_1(t)| &\le \sum_{i,j=1}^{2}\int_{\mathbb{R}^3}|\omega_i||\partial_i u_j||\omega_j|dx \\ &\le \sum_{i,j=1}^{2}\|\partial_i u_j\|_{L^\alpha}\|\omega_i\|_{L^p}\|\omega_j\|_{L^p} \\ &\le \sum_{i,j=1}^{2}\|\partial_i u_j\|_{L^\alpha}\|\omega\|_{L^p}^2, \end{aligned} \tag{2.4}$$

where α and p satisfy

$$\frac{1}{\alpha}+\frac{2}{p}=1 \text{ with } \alpha>3,\ 2\le p<3. \tag{2.5}$$

By using of (1.36), (2.4) follows that

$$|I_1(t)|\le C\|\partial_3 u_3\|_{L^\alpha}\|\omega\|_{L^p}^2+C\|\omega_3\|_{L^\alpha}\|\omega\|_{L^p}^2. \tag{2.6}$$

For $I_2(t)$, by virtue of (1.31), (2.5) and Hölder's inequality, one has

$$\begin{aligned} |I_2(t)| &\le \sum_{i=1}^{2}\int_{\mathbb{R}^3}|\omega_i||\partial_i u_3||\omega_3|dx \\ &\le \sum_{i=1}^{2}\|\omega_3\|_{L^\alpha}\|\partial_i u_3\|_{L^p}\|\omega_i\|_{L^p} \\ &\le C\|\omega_3\|_{L^\alpha}\|\nabla u\|_{L^p}\|\omega\|_{L^p} \\ &\le C\|\omega_3\|_{L^\alpha}\|\omega\|_{L^p}^2. \end{aligned} \tag{2.7}$$

Similar to $I_2(t)$, we have the estimates for $I_3(t)$

$$|I_3(t)|\le C\|\omega_3\|_{L^\alpha}\|\omega\|_{L^p}^2. \tag{2.8}$$

Combining (2.5), (2.7) and (2.8), we have

$$\frac{1}{2}\frac{d}{dt}\|\omega\|_{L^2}^2+\nu\|\nabla\omega\|_{L^2}^2\le C\|\partial_3 u_3\|_{L^\alpha}\|\omega\|_{L^p}^2+C\|\omega_3\|_{L^\alpha}\|\omega\|_{L^p}^2. \tag{2.9}$$

Applying (1.28) and Young's inequality, it follows that

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|\omega\|_{L^2}^2+\nu\|\nabla\omega\|_{L^2}^2 &\le C\|\partial_3 u_3\|_{L^\alpha}\|\omega\|_{L^p}^2+C\|\omega_3\|_{L^\alpha}\|\omega\|_{L^p}^2 \\ &\le C\|\partial_3 u_3\|_{L^\alpha}\|\omega\|_{L^2}^{\frac{6-p}{p}}\|\nabla\omega\|_{L^2}^{\frac{3p-6}{p}} \\ &\quad +C\|\omega_3\|_{L^\alpha}\|\omega\|_{L^2}^{\frac{6-p}{p}}\|\nabla\omega\|_{L^2}^{\frac{3p-6}{p}} \\ &\le C\|\partial_3 u_3\|_{L^\alpha}^{\frac{2p}{6-p}}\|\omega\|_{L^2}^2 \\ &\quad +C\|\omega_3\|_{L^\alpha}^{\frac{2p}{6-p}}\|\omega\|_{L^2}^2+\frac{\nu}{2}\|\nabla\omega\|_{L^2}^2 \\ &= C\|\partial_3 u_3\|_{L^\alpha}^{\frac{2\alpha}{2\alpha-3}}\|\omega\|_{L^2}^2 \\ &\quad +C\|\omega_3\|_{L^\alpha}^{\frac{2\alpha}{2\alpha-3}}\|\omega\|_{L^2}^2+\frac{\nu}{2}\|\nabla\omega\|_{L^2}^2. \end{aligned} \tag{2.10}$$

Integrating (2.10) and absorbing the last term, one has

$$\begin{aligned}\|\omega\|_{L^2}^2 &+ \nu\int_0^t \|\nabla\omega\|_{L^2}^2 d\tau \\ &\le C\int_0^t \|\partial_3 u_3\|_{L^\alpha}^{\frac{2\alpha}{2\alpha-3}}\|\omega\|_{L^2}^2 d\tau + C\int_0^t \|\omega_3\|_{L^\alpha}^{\frac{2\alpha}{2\alpha-3}}\|\omega\|_{L^2}^2 d\tau + \|\omega(0)\|_{L^2}^2.\end{aligned} \tag{2.11}$$

By using of Gronwall's inequality, we obtain

$$\begin{aligned}\|\omega\|_{L^2}^2 &+ \nu\int_0^t \|\nabla\omega\|_{L^2}^2 d\tau \\ &\le \left(\|\omega(0)\|_{L^2}^2\right)\exp\left(C\int_0^t \|\partial_3 u_3\|_{L^\alpha}^{\frac{2\alpha}{2\alpha-3}} d\tau\right)\exp\left(C\int_0^t \|\omega_3\|_{L^\alpha}^{\frac{2\alpha}{2\alpha-3}} d\tau\right),\end{aligned} \tag{2.12}$$

by the condition (2.3) and (2.3), we have

$$\omega \in L^\infty(0,T^*;L^2(\mathbb{R}^3)) \cap L^2(0,T^*;H^1(\mathbb{R}^3)).$$

Therefore, by (1.31) and (1.35), we get that the H^1 norm of the strong solution u is bounded on the maximal interval of existence $(0,T^*)$. The proof is completed.
Proof of Theorem 1.3 Taking the inner product of the equation (1.1) with $-\Delta u$ in L^2, we obtain

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\|\nabla u\|_2^2 \quad &+\nu\|\Delta u\|_2^2 \\ &= \sum_{i,j,k=1}^{3}\int_{\mathbb{R}^3} u_i\partial_i u_j \partial_{kk} u_j dx \\ &= \sum_{i,j,k=1}^{2}\int_{\mathbb{R}^3} u_i\partial_i u_j \partial_{kk} u_j dx + \sum_{i=1}^{3}\sum_{j=1}^{2}\int_{\mathbb{R}^3} u_i\partial_i u_j \partial_{33} u_j dx \\ &\quad + \sum_{j=1}^{2}\sum_{k=1}^{2}\int_{\mathbb{R}^3} u_3\partial_3 u_j \partial_{kk} u_j dx + \sum_{i,k=1}^{3}\int_{\mathbb{R}^3} u_i\partial_i u_3 \partial_{kk} u_3 dx \\ &= I_1(t) + I_2(t) + I_3(t) + I_4(t).\end{aligned} \tag{2.13}$$

By integrating by parts a few times and using the incompressibility condition, we have

$$I_1(t) = \frac{1}{2}\sum_{i,j,k=1}^{2}\int_{\mathbb{R}^3} \partial_i u_i \partial_k u_j \partial_k u_j dx - \sum_{i,j,k=1}^{2}\int_{\mathbb{R}^3} \partial_k u_i \partial_i u_j \partial_k u_j dx = I_1^1(t) + I_1^2(t).$$

The terms $I_1^1(t)$, $I_1^2(t)$, $I_3(t)$ and $I_4(t)$ read as

$$I_1^1(t) = -\frac{1}{2}\sum_{j,k=1}^{2}\int_{\mathbb{R}^3} \partial_3 u_3 \partial_k u_j \partial_k u_j dx,$$

$$
\begin{aligned}
I_1^2(t) &= -\sum_{i,j,k=1}^{2} \int_{\mathbb{R}^3} \partial_k u_i \partial_i u_j \partial_k u_j dx \\
&= \int_{\mathbb{R}^3} \partial_2 u_1 \partial_1 u_2 \partial_2 u_2 dx + \int_{\mathbb{R}^3} \partial_1 u_2 \partial_2 u_1 \partial_1 u_1 dx + \int_{\mathbb{R}^3} \partial_1 u_1 \partial_1 u_2 \partial_1 u_2 dx \\
&\quad + \int_{\mathbb{R}^3} \partial_1 u_2 \partial_2 u_2 \partial_1 u_2 dx + \int_{\mathbb{R}^3} \partial_2 u_1 \partial_1 u_1 \partial_2 u_1 dx + \int_{\mathbb{R}^3} \partial_1 u_1 \partial_1 u_1 \partial_1 u_1 dx \\
&\quad + \int_{\mathbb{R}^3} \partial_2 u_2 \partial_2 u_2 \partial_2 u_2 dx + \int_{\mathbb{R}^3} \partial_2 u_2 \partial_2 u_1 \partial_2 u_1 dx \\
&= -\int_{\mathbb{R}^3} (\partial_2 u_1 \partial_1 u_2 \partial_3 u_3 + \partial_3 u_3 \partial_1 u_2 \partial_1 u_2 + \partial_2 u_1 \partial_3 u_3 \partial_2 u_1) dx \\
&\quad - \int_{\mathbb{R}^3} (\partial_1 u_1 \partial_1 u_1 \partial_3 u_3 + \partial_3 u_3 \partial_2 u_2 \partial_2 u_2 - \partial_1 u_1 \partial_3 u_3 \partial_2 u_2) dx, \\
I_3(t) &= \sum_{j=1}^{2} \sum_{k=1}^{2} \int_{\mathbb{R}^3} u_3 \partial_3 u_j \partial_{kk} u_j dx \\
&= -\sum_{j=1}^{2} \sum_{k=1}^{2} \int_{\mathbb{R}^3} \partial_k u_3 \partial_3 u_j \partial_k u_j dx - \sum_{j=1}^{2} \sum_{k=1}^{2} \int_{\mathbb{R}^3} u_3 \partial_{3k} u_j \partial_k u_j dx \\
&= -\sum_{j=1}^{2} \sum_{k=1}^{2} \int_{\mathbb{R}^3} \partial_k u_3 \partial_3 u_j \partial_k u_j dx + \frac{1}{2} \sum_{j=1}^{2} \sum_{k=1}^{2} \int_{\mathbb{R}^3} \partial_3 u_3 \partial_k u_j \partial_k u_j dx, \\
I_4(t) &= \sum_{i,k=1}^{3} \int_{\mathbb{R}^3} u_i \partial_i u_3 \partial_{kk} u_3 dx \\
&= -\sum_{i,k=1}^{3} \int_{\mathbb{R}^3} \partial_k u_i \partial_i u_3 \partial_k u_3 dx - \sum_{i,k=1}^{3} \int_{\mathbb{R}^3} u_i \partial_{ik} u_3 \partial_k u_3 dx \\
&= -\sum_{i,k=1}^{3} \int_{\mathbb{R}^3} \partial_k u_i \partial_i u_3 \partial_k u_3 dx.
\end{aligned}
$$

The above four equalities imply that

$$|I_i| \le C \int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx, \ i = 1, 3, 4.$$

- Case of $\partial_3 u_2, \partial_3 u_3$

As for I_2, we have

$$
\begin{aligned}
I_2 &= -\int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_1 \partial_3 u_1 + \partial_3 u_2 \partial_2 u_1 \partial_3 u_1 + \partial_3 u_3 \partial_3 u_1 \partial_3 u_1 dx \\
&\qquad - \int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_2 \partial_3 u_2 + \partial_3 u_2 \partial_2 u_2 \partial_3 u_2 + \partial_3 u_3 \partial_3 u_2 \partial_3 u_2 dx \\
&= I_2^1 + I_2^2 + I_2^3 + I_2^4 + I_2^5 + I_2^6.
\end{aligned}
$$

It is obvious that

$$|I_2^j| \le C \int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx, \ j = 3, 6, \tag{2.14}$$

and

$$|I_2^j| \le C \int_{\mathbb{R}^3} |\partial_3 u_2||\nabla u|^2 dx, \ j = 2, 4, 5. \tag{2.15}$$

For the first term, by using the incompressibility condition, and integrating by parts a few times, we note that

$$\begin{aligned} -\int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_1 \partial_3 u_1 dx &= 2\int_{\mathbb{R}^3} \partial_{13} u_1 \partial_3 u_1 u_1 dx \\ &= -2\int_{\mathbb{R}^3} \partial_{23} u_2 \partial_3 u_1 u_1 dx - 2\int_{\mathbb{R}^3} \partial_{33} u_3 \partial_3 u_1 u_1 dx \\ &= 2\int_{\mathbb{R}^3} \partial_{23} u_1 \partial_3 u_2 u_1 dx + 2\int_{\mathbb{R}^3} \partial_3 u_2 \partial_3 u_1 \partial_2 u_1 dx \\ &\quad +2\int_{\mathbb{R}^3} \partial_{33} u_1 \partial_3 u_3 u_1 dx + 2\int_{\mathbb{R}^3} \partial_3 u_3 \partial_3 u_1 \partial_3 u_1 dx \end{aligned} \tag{2.16}$$

As before, I_2^1 has the estimate

$$\begin{aligned} |I_2^1| &\le C\int_{\mathbb{R}^3} |\partial_3 u_2||\nabla u|^2 dx + C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx \\ &\quad +C\int_{\mathbb{R}^3} |\partial_3 u_2||\nabla^2 u||u| dx + C\int_{\mathbb{R}^3} |\partial_3 u_3||\nabla^2 u||u| dx. \end{aligned} \tag{2.17}$$

Therefore, by above inequalities, we see that

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|\nabla u\|_2^2 + \nu\|\Delta u\|_2^2 &\le C\int_{\mathbb{R}^3} |\partial_3 u_2||\nabla u|^2 dx + C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx \\ &\quad +C\int_{\mathbb{R}^3} |\partial_3 u_2||\nabla^2 u||u| dx + C\int_{\mathbb{R}^3} |\partial_3 u_3||\nabla^2 u||u| dx. \end{aligned} \tag{2.18}$$

Let $T' \in (0,T)$ with $T' \le T^*$ be arbitrary. We shall prove that T' is not a blow-up point. By decreasing of α if necessary, we may assume

$$u_3 \in L^{\beta_1}(0,T;L^{\alpha_1}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_1} + \frac{2}{\beta_1} = 1, \alpha_1 \in (3,\infty], \beta_1 \in [2,\infty), \tag{2.19}$$

and

$$\partial_3 u_i \in L^{\beta_2}(0,T;L^{\alpha_2}(\mathbb{R}^3)) \text{ with } \frac{3}{\alpha_2} + \frac{2}{\beta_2} = 2, \alpha_2 \in [2,3], \beta_2 \in [2,\infty), i = 2,3. \tag{2.20}$$

Choose a $t_1 \in (0,T')$ such that

$$\|\partial_3 u_i\|_{L^{\beta_2}(t_1,T';L^{\alpha_2}(\mathbb{R}^3))} \le \epsilon, \;\; i = 2,3, \tag{2.21}$$

and

$$\|u_3\|_{L^{\beta_1}(t_1,T';L^{\alpha_1}(\mathbb{R}^3))} \le \epsilon, \tag{2.22}$$

where ϵ is to be determined. Let $t_2 \in (t_1,T')$ be arbitrary. For any r, s, we abbreviate

$$\|\cdot\|_{L_t^s L_x^r} = \|\cdot\|_{L^s((t_1,t_2);L^r(\mathbb{R}^3))}. \tag{2.23}$$

We denote

$$L = \|\nabla u\|_{L_t^\infty L_x^2} + \nu\|\Delta u\|_{L_t^2 L_x^2}. \tag{2.24}$$

Let $t \in (t_1, t_2]$ be arbitrary. integrating (2.18) on (t_1, t), we get

$$\begin{aligned}\|\nabla u\|_{L^2}^2 \quad & +2\nu\int_{t_1}^{t}\|\Delta u\|_{L^2}^2 d\tau \\ & \leq C\int_{t_1}^{t}\int_{\mathbb{R}^3}|\partial_3 u_2||\nabla u|^2 dxd\tau + C\int_{t_1}^{t}\int_{\mathbb{R}^3}|u_3||\nabla u||\nabla^2 u|dxd\tau \\ & \quad +C\int_{t_1}^{t}\int_{\mathbb{R}^3}|\partial_3 u_2||\nabla^2 u||u|dxd\tau + C\int_{t_1}^{t}\int_{\mathbb{R}^3}|\partial_3 u_3||\nabla^2 u||u|dxd\tau \\ & \quad +\|\nabla u(t_1)\|_{L^2}^2 \\ & = L_1 + L_2 + L_3 + L_4 + L_5.\end{aligned} \tag{2.25}$$

We estimate L_i, $i = 1, 2, 3, 4$ one by one. Firstly, for L_1 we have

$$L_1 \quad \leq C\|\partial_3 u_2\|_{L_t^{\beta_2}L_x^{\alpha_2}}\|\nabla u\|_{L_t^{s_1}L_x^{r_1}}^2 \tag{2.26}$$

where s_1 and r_1 satisfy

$$\frac{1}{\alpha_2} + \frac{2}{r_1} = 1, \quad \frac{1}{\beta_2} + \frac{2}{s_1} = 1, \tag{2.27}$$

by (2.20) and (2.27), we have $3 \leq r_1 \leq 4$ and $2/s_1 + 3/r_1 = 3/2$. By Lemma 2.1, we have

$$L_1 \leq C\epsilon L^2. \tag{2.28}$$

As for L_2, we have

$$\begin{aligned}L_2 \quad & \leq C\|u_3\|_{L_t^{\beta_1}L_x^{\alpha_1}}\|\Delta u\|_{L_t^2 L_x^2}\|\nabla u\|_{L_t^{s_2}L_x^{r_2}} \\ & \leq C\epsilon L^2,\end{aligned} \tag{2.29}$$

where s_2 and r_2 satisfy

$$\frac{1}{\alpha_1} + \frac{1}{r_2} = \frac{1}{2}, \quad \frac{1}{\beta_1} + \frac{1}{s_2} = \frac{1}{2}, \tag{2.30}$$

by (2.19) and (2.30), we have $2 \leq r_2 < 6$ and $2/s_2 + 3/r_2 = 3/2$. thus, s_2 and r_2 satisfy Lemma 2.1. The estimate of L_3 is as follows

$$L_3 \quad \leq C\|\partial_3 u_2\|_{L_t^{\beta_2}L_x^{\alpha_2}}\|\Delta u\|_{L_t^2 L_x^2}\|u\|_{L_t^{s_3}L_x^{r_3}} \tag{2.31}$$

where s_3 and r_3 satisfy

$$\frac{1}{\alpha_2} + \frac{1}{r_3} = \frac{1}{2}, \quad \frac{1}{\beta_2} + \frac{1}{s_3} = \frac{1}{2}, \tag{2.32}$$

by (2.20) and (2.32), we have

$$\frac{2}{s_3} + \frac{3}{r_3} = \frac{1}{2}, \ r_3 \geq 6. \tag{2.33}$$

By the Gagliardo-Nirenberg inequality

$$\|v\|_{L^r} \leq \|\nabla v\|_{L^{\frac{3r}{3+r}}}, \ \frac{3}{2} \leq r < \infty. \tag{2.34}$$

Therefore, we have

$$\|u\|_{L_t^{s_3}L_x^{r_3}} \leq \|\nabla u\|_{L_t^{s_3}L_x^{r_4}}, \tag{2.35}$$

where $r_4 = 3r_3/(3 + r_3)$, and we have $2/s_3 + 3/r_4 = 3/2$ with $2 \leq r_4 \leq 6$. Combining (2.20), (2.31) and (2.35), as well as Lemma 2.1, on has

$$L_3 \leq C\epsilon L^2. \tag{2.36}$$

The term L_4 is estimated the same way and we get the same result. Finally, we obtain

$$L^2 \leq C\epsilon L^2 + \|\nabla u(t_1)\|_{L^2}^2.$$

If t_1 is sufficiently close to T' and ϵ is sufficiently small, we can absorb the first term into the left hand side, and then we obtain that L is bounded with a bound independent of $t_2 \in (t_1, T')$. Finally, we get $\|\nabla u\|_{L_t^\infty L_x^2(\mathbb{R}^3\times(t_1,T'))} \leq C$. Therefore, the solution cannot blow up at T'. We complete the proof of the case of $\partial_3 u_2, \partial_3 u_3$.

- Case of $\partial_3 u_1, \partial_3 u_3$

This case is similar to the first case. The main difference is that we have another estimate for L_2. Also by using the incompressibility condition, and integrating by parts a few times, we note that I_2^5 becomes

$$\begin{aligned}
-\int_{\mathbb{R}^3} \partial_3 u_2 \partial_3 u_2 \partial_2 u_2 dx &= 2\int_{\mathbb{R}^3} \partial_{23} u_2 \partial_3 u_2 u_2 dx \\
&= -2\int_{\mathbb{R}^3} \partial_{13} u_1 \partial_3 u_2 u_2 dx - 2\int_{\mathbb{R}^3} \partial_{33} u_3 \partial_3 u_2 u_2 dx \\
&= 2\int_{\mathbb{R}^3} \partial_{13} u_2 \partial_3 u_1 u_2 dx + 2\int_{\mathbb{R}^3} \partial_3 u_2 \partial_3 u_1 \partial_1 u_2 dx \\
&\quad +2\int_{\mathbb{R}^3} \partial_{33} u_2 \partial_3 u_3 u_2 dx - 2\int_{\mathbb{R}^3} \partial_3 u_3 \partial_3 u_2 \partial_3 u_2 dx,
\end{aligned} \tag{2.37}$$

and then I_2^5 has the estimate

$$\begin{aligned}
|I_2^5| &\leq C\int_{\mathbb{R}^3} |\partial_3 u_1||\nabla u|^2 dx + C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx \\
&\quad +C\int_{\mathbb{R}^3} |\partial_3 u_1||\nabla^2 u||u| dx + C\int_{\mathbb{R}^3} |\partial_3 u_3||\nabla^2 u||u| dx.
\end{aligned} \tag{2.38}$$

Therefore, we finally get

$$\begin{aligned}
\frac{1}{2}\frac{d}{dt}\|\nabla u\|_2^2 + \nu\|\Delta u\|_2^2 &\leq C\int_{\mathbb{R}^3} |\partial_3 u_1||\nabla u|^2 dx + C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx \\
&\quad +C\int_{\mathbb{R}^3} |\partial_3 u_1||\nabla^2 u||u| dx + C\int_{\mathbb{R}^3} |\partial_3 u_3||\nabla^2 u||u| dx.
\end{aligned} \tag{2.39}$$

By using (2.39), we give the same method as before to get the desired result. The proof is completed.

Proof of Theorem 1.4 From the condition of this Theorem, we split the proof into two parts.

- $i \neq j$

For convenience of writing, for every $\alpha_2 > 2$, we set

$$r = \frac{3\alpha_2 - 2}{\alpha_2}. \tag{2.40}$$

It is easy to check that $r > 2$ when $\alpha_2 > 2$. Without loss of generality, in the proof, we will assume that $i = 1, j = 2$, the other cases can be discussed in the same way (for details see Remark 2.2 below). We begin with (2.13), and the same process to the proof of Theorem 1.3, we firstly have

$$|I_i| \leq C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx,\ i = 1, 3, 4.$$

As for I_2, also by the incompressibility condition, we have

$$\begin{aligned} I_2 &= -\int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_1 \partial_3 u_1 + \partial_3 u_2 \partial_2 u_1 \partial_3 u_1 + \partial_3 u_3 \partial_3 u_1 \partial_3 u_1 dx \\ &\quad - \int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_2 \partial_3 u_2 + \partial_3 u_2 \partial_2 u_2 \partial_3 u_2 + \partial_3 u_3 \partial_3 u_2 \partial_3 u_2 dx \\ &= -\int_{\mathbb{R}^3} \partial_3 u_1 \left(-\partial_2 u_2 - \partial_3 u_3\right) \partial_3 u_1 + \partial_3 u_2 \partial_2 u_1 \partial_3 u_1 + \partial_3 u_3 \partial_3 u_1 \partial_3 u_1 dx \\ &\quad - \int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_2 \partial_3 u_2 + \partial_3 u_2 \partial_2 u_2 \partial_3 u_2 + \partial_3 u_3 \partial_3 u_2 \partial_3 u_2 dx \\ &\leq \int_{\mathbb{R}^3} |u_2||\nabla u||\nabla^2 u| dx + \int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx. \end{aligned}$$

Therefore, we get

$$\begin{aligned} &\frac{1}{2}\frac{d}{dt}\|\nabla u\|_{L^2}^2 + \nu\|\Delta u\|_{L^2}^2 \\ &\qquad \leq C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx + C\int_{\mathbb{R}^3} |u_2||\nabla u||\nabla^2 u| dx, \\ &\qquad = K_1(t) + K_2(t). \end{aligned} \tag{2.41}$$

Next, we estimate $K_1(t)$ and $K_2(t)$. Firstly, we pay attention to $K_2(t)$, applying Hölder's inequality several times, we obtain

$$\begin{aligned} K_2(t) &\leq C\int_{\mathbb{R}^2} \max_{x_1} |u_2| \left(\int_{\mathbb{R}} |\nabla u|^2 dx_1\right)^{\frac{1}{2}} \left(\int_{\mathbb{R}} |\nabla^2 u|^2 dx_1\right)^{\frac{1}{2}} dx_2 dx_3 \\ &\leq C\Big[\int_{\mathbb{R}^2} (\max_{x_1} |u_2|)^r dx_2 dx_3\Big]^{\frac{1}{r}} \Big[\int_{\mathbb{R}^2} \Big(\int_{\mathbb{R}} |\nabla u|^2 dx_1\Big)^{\frac{r}{r-2}} dx_2 dx_3\Big]^{\frac{r-2}{2r}} \\ &\quad \times\Big[\int_{\mathbb{R}^3} |\nabla^2 u|^2 dx_1 dx_2 dx_3\Big]^{\frac{1}{2}} \\ &\leq C\Big[\int_{\mathbb{R}^3} |u_2|^{r-1} \partial_1 u_2 dx_1 dx_2 dx_3\Big]^{\frac{1}{r}} \|\Delta u\|_2 \\ &\quad \times\Big[\int_{\mathbb{R}} \Big(\int_{\mathbb{R}^2} |\nabla u|^{\frac{2r}{r-2}} dx_2 dx_3\Big)^{\frac{r-2}{r}} dx_1\Big]^{\frac{1}{2}} \\ &\leq C\|u_2\|_{L^2}^{\frac{r-1}{r}} \|\partial_1 u_2\|_{L^{\frac{2}{3-r}}}^{\frac{1}{r}} \|\nabla u\|_{L^2}^{\frac{r-2}{r}} \|\partial_2 \nabla u\|_{L^2}^{\frac{1}{r}} \|\partial_3 \nabla u\|_{L^2}^{\frac{1}{r}} \|\Delta u\|_2 \\ &\leq C\|u_2\|_{L^2}^{\frac{r-1}{r}} \|\partial_1 u_2\|_{L^{\frac{2}{3-r}}}^{\frac{1}{r}} \|\nabla u\|_{L^2}^{\frac{r-2}{r}} \|\Delta u\|_{L^2}^{\frac{r+2}{r}}. \end{aligned} \tag{2.42}$$

In above inequality, from (2.40), we note that $\frac{2}{3-r} = \alpha_2$. Therefore, applying Young's inequality, (2.42) immediately implies

$$\begin{aligned} K_2(t) &\leq C\|u_2\|_{L^2}^{\frac{2(\alpha_2-1)}{3\alpha_2-2}} \|\partial_1 u_2\|_{L^{\alpha_2}}^{\frac{\alpha_2}{3\alpha_2-2}} \|\nabla u\|_{L^2}^{\frac{\alpha_2-2}{3\alpha_2-2}} \|\Delta u\|_{L^2}^{\frac{5\alpha_2-2}{3\alpha_2-2}} \\ &\leq C\|u_2\|_{L^2}^{\frac{4(\alpha_2-1)}{\alpha_2-2}} \|\partial_1 u_2\|_{L^{\alpha_2}}^{\frac{2\alpha_2}{\alpha_2-2}} \|\nabla u\|_{L^2}^2 + \frac{\nu}{4}\|\Delta u\|_{L^2}^2. \end{aligned} \tag{2.43}$$

As for $K_1(t)$, applying Hölder's and Young's inequalities, we have

$$\begin{aligned} K_1(t) &= C\int_{\mathbb{R}^3}|u_3||\nabla u||\nabla^2 u|dx \\ &\leq C\|u_3\|_{L^{\alpha_1}}\|\nabla u\|_{L^q}\|\Delta u\|_{L^2} \\ &\leq C\|u_3\|_{L^{\alpha_1}}\|\nabla u\|_{L^2}^{\frac{6-q}{2q}}\|\Delta u\|_{L^2}^{\frac{5q-6}{2q}} \\ &\leq C\|u_3\|_{L^{\alpha_1}}^{\frac{4q}{6-q}}\|\nabla u\|_{L^2}^2+\frac{\nu}{4}\|\Delta u\|_{L^2}^2 \end{aligned} \tag{2.44}$$

where α_1 and q satisfy

$$\frac{1}{q}+\frac{1}{\alpha_1}=\frac{1}{2} \text{ with } 2\leq q<6, \alpha_1>3. \tag{2.45}$$

From (2.44) and (2.43), one has

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|\nabla u\|_{L^2}^2+\nu\|\Delta u\|_{L^2}^2 &\leq C\|u_2\|_{L^2}^{\frac{4(\alpha-1)}{\alpha-2}}\|\partial_1 u_2\|_{L^\alpha}^{\frac{2\alpha}{\alpha-2}}\|\nabla u\|_{L^2}^2 \\ &\quad +C\|u_3\|_{L^\alpha}^{\frac{2\alpha}{\alpha-3}}\|\nabla u\|_{L^2}^2+\frac{\nu}{2}\|\Delta u\|_{L^2}^2. \end{aligned} \tag{2.46}$$

Absorbing the last term in right hand of (2.46) and integrating the inequality on time, using the energy inequality, we obtain

$$\begin{aligned} \|\nabla u\|_{L^2}^2+\nu\int_0^t\|\nabla\Delta u\|_{L^2}^2d\tau &\leq C\int_0^t\|u_2\|_{L^2}^{\frac{4(\alpha_2-1)}{\alpha_2-2}}\|\partial_1 u_2\|_{L^{\alpha_2}}^{\frac{2\alpha_2}{\alpha_2-2}}\|\nabla u\|_{L^2}^2d\tau \\ &\quad +C\int_0^t\|u_3\|_{L^{\alpha_1}}^{\frac{2\alpha_1}{\alpha_1-3}}\|\nabla u\|_{L^2}^2d\tau+\|\nabla u(0)\|_{L^2}^2 \\ &\leq C\int_0^t\|\partial_1 u_2\|_{L^{\alpha_2}}^{\frac{2\alpha_2}{\alpha_2-2}}\|\nabla u\|_{L^2}^2d\tau \\ &\quad +C\int_0^t\|u_3\|_{L^{\alpha_1}}^{\frac{2\alpha_1}{\alpha_1-3}}\|\nabla u\|_{L^2}^2d\tau+\|\nabla u(0)\|_{L^2}^2. \end{aligned} \tag{2.47}$$

By using Gronwall's inequality, we obtain

$$\begin{aligned} &\|\nabla u\|_{L^2}^2+\nu\int_0^t\|\Delta u\|_{L^2}^2d\tau \\ &\quad\leq\left(\|\nabla u(0)\|_{L^2}^2\right)\exp\left(C\int_0^t\|\partial_1 u_2\|_{L^{\alpha_2}}^{\frac{2\alpha_2}{\alpha_2-2}}d\tau\right)\exp\left(C\int_0^t\|u_3\|_{L^{\alpha_1}}^{\frac{2\alpha_1}{\alpha_1-3}}d\tau\right) \end{aligned} \tag{2.48}$$

By condition (1.11)−(1.13), (2.48) follows that the H^1 norm of the strong solution u is bounded on the maximal interval of existence $(0,T^*)$. Thus we prove (a).

- $i=j$

Without loss of generality, here, we assume $i=j=2$. Similar to the proof of the part (a), we estimate the second term $K_2(t)$ of (2.41). Firstly, for every

$$\alpha_3\geq 2, \tag{2.49}$$

we choose

$$\begin{cases} \dfrac{1}{\gamma} = \dfrac{3+\sqrt{24\alpha_3^2-24\alpha_3+9}}{12\alpha_3}, \\ \mu = \dfrac{\alpha_3\gamma}{2\alpha_3-\gamma}. \end{cases} \tag{2.50}$$

From $(2.50)_1$ and (2.49), we have $\frac{1}{\gamma} > \frac{1}{2\alpha_3}$, which means that $\gamma < 2\alpha_3$, and hance $\mu > 0$ is well defined in $(2.50)_2$. Also from $(2.50)_1$, we see that γ is an increasing function with the variable α_3, and from (2.49) we get

$$2 < \frac{9}{4} < \delta \leq \gamma < \sqrt{6}, \tag{2.51}$$

where $\delta = \frac{-3+\sqrt{57}}{2}$, and besides (2.50) follows

$$\frac{1}{\mu} + \frac{1}{\alpha_3} + \frac{\gamma-2}{\gamma} = 1. \tag{2.52}$$

Moreover, from the fact $\gamma > 2$ and $(2.50)_2$, it is easy to check that $\mu > 1$. On the other hand, we see that γ satisfies

$$\alpha_3 = \frac{3\gamma-\gamma^2}{6-\gamma^2}, \tag{2.53}$$

and

$$\frac{3\gamma-\gamma^2}{6-\gamma^2} - \frac{3\gamma}{6-\gamma} = \frac{\gamma^2(4\gamma-9)}{(6-\gamma^2)(6-\gamma)} > 0 \text{ with } \delta \leq \gamma < \sqrt{6}.$$

Combining (2.53) and above inequality, γ satisfies $\alpha_2 > \frac{3\gamma}{6-\gamma}$, and hance by the definition of μ in (2.50), a short calculation shows that $\mu < 3$. Finally, we obtain

$$1 < \mu < 3. \tag{2.54}$$

We choose

$$\beta_3 = \frac{2\mu}{3-\mu}, \tag{2.55}$$

then we have

$$\frac{1}{\mu} = \frac{2}{3}\left(\frac{1}{\beta_3} + \frac{1}{2}\right),$$

by $(2.50)_1$, (2.52) and (2.55), we can compute that

$$\frac{2}{\beta_3} + 1 = \frac{3}{\mu} = \frac{6}{\gamma} - \frac{3}{\alpha_3} = \frac{-3+\sqrt{24\alpha_3^2-24\alpha_3+9}}{2\alpha_3} \Rightarrow \frac{3}{2\alpha_3} + \frac{2}{\beta_3} = f(\alpha_3). \tag{2.56}$$

Now, we use $|u_2|^{\gamma-2}u_2$ with $\gamma > 2$ as test function in the equation (1.1) for u_2. By using of Gagliardo-Nirenberg and Hölder's inequalities, we have

$$\begin{aligned}\frac{1}{\gamma}\frac{d}{dt}\|u_2\|_\gamma^\gamma \quad & +C(\gamma)\nu\|\nabla|u_2|^{\frac{\gamma}{2}}\|_{L^2}^2 \\ & = -\int_{\mathbb{R}^3}\partial_2 p|u_2|^{\gamma-2}u_2dx \\ & \le C\int_{\mathbb{R}^3}|p||u_2|^{\gamma-2}|\partial_2 u_2|dx \\ & \le C\|p\|_{L^\mu}\|u_2\|_{L^\gamma}^{\gamma-2}\|\partial_2 u_2\|_{L^{\alpha_3}} \\ & \le C\|u\|_{L^{2\mu}}^2\|u_2\|_{L^\gamma}^{\gamma-2}\|\partial_2 u_2\|_{L^{\alpha_3}} \text{ (by (1.30))} \\ & \le C\|u\|_{L^2}^{\frac{3-\mu}{\mu}}\|\nabla u\|_{L^2}^{\frac{3(\mu-1)}{\mu}}\|u_2\|_{L^\gamma}^{\gamma-2}\|\partial_2 u_2\|_{L^{\alpha_3}},\end{aligned}\tag{2.57}$$

where the γ, μ and α_3 satisfy (2.52). The above inequality immediately implies that

$$\frac{1}{2}\frac{d}{dt}\|u_2\|_{L^\gamma}^2 \le C\|u\|_{L^2}^{\frac{3-\mu}{\mu}}\|\nabla u\|_{L^2}^{\frac{3(\mu-1)}{\mu}}\|\partial_2 u_2\|_{L^{\alpha_3}}.\tag{2.58}$$

In view of (2.54), we have $\frac{3(\mu-1)}{\mu} < 2$, applying Young's inequality, we have

$$\frac{1}{2}\frac{d}{dt}\|u_2\|_{L^\gamma}^2 \le C\|\nabla u\|_{L^2}^2 + \|u\|_{L^2}^2\|\partial_2 u_2\|_{L^{\alpha_3}}^{\frac{2\mu}{3-\mu}}.\tag{2.59}$$

Integrating (2.59) on time, and by energy inequality (2.1), we obtain

$$\begin{aligned}\|u_2\|_{L^\gamma}^2 \quad & \le \|u_2(0)\|_{L^\gamma}^2 + C + C\int_0^t \|\partial_2 u_2\|_{L^{\alpha_3}}^{\frac{2\mu}{3-\mu}}d\tau \\ & = \|u_2(0)\|_{L^\gamma}^2 + C + C\int_0^t \|\partial_2 u_2\|_{L^{\alpha_3}}^{\beta_3}d\tau.\end{aligned}\tag{2.60}$$

In view of (2.51), we have $\|u_2(0)\|_\gamma < C$ for some $C > 0$, therefore, by the condition (1.15), we get

$$u_2 \in L^\infty(0,T;L^\gamma(\mathbb{R}^3)).\tag{2.61}$$

For the mentioned parameter γ in (2.50), we set

$$r = \frac{(\gamma+1)\alpha_3 - \gamma}{\alpha_3},\tag{2.62}$$

since $\alpha_3 \ge 2$ and $\gamma > \delta$, it is easy to see $r > 2$. Similar to (2.42), also by Hölder's and Young's inequalities, we obtain another estimate

$$\begin{aligned}K_2(t) \quad & \le C\|u_2\|_{L^\gamma}^{\frac{r-1}{r}}\|\partial_2 u_2\|_{L^{\frac{\gamma}{\gamma-r+1}}}^{\frac{1}{r}}\|\nabla u\|_{L^2}^{\frac{r-2}{r}}\|\Delta u\|_{L^2}^{\frac{r+2}{r}} \\ & \le C\|u_2\|_{L^\gamma}^{\frac{2(r-1)}{r-2}}\|\partial_2 u_2\|_{L^{\alpha_3}}^{\frac{2}{r-2}}\|\nabla u\|_{L^2}^2 + \frac{\nu}{4}\|\Delta u\|_{L^2}^2.\end{aligned}\tag{2.63}$$

Applying (2.61) and integrating above inequality, one has

$$\begin{aligned}\int_0^t K_2(\tau)d\tau &\le C\int_0^t \|u_2\|_{L^\gamma}^{\frac{2(r-1)}{r-2}}\|\partial_2 u_2\|_{L^{\alpha_3}}^{\frac{2}{r-2}}\|\nabla u\|_{L^2}^2 d\tau+\frac{\nu}{4}\int_0^t\|\Delta u\|_{L^2}^2 d\tau\\ &\le C\int_0^t \|\partial_2 u_2\|_{L^{\alpha_3}}^{\frac{2}{r-2}}\|\nabla u\|_{L^2}^2 d\tau+\frac{\nu}{4}\int_0^t\|\Delta u\|_{L^2}^2 d\tau.\end{aligned} \tag{2.64}$$

Integrating (2.41) on time, and absorbing the last term in (2.64) and (2.44) respectively, it follows that

$$\begin{aligned}&\|\nabla u\|_{L^2}^2+\nu\int_0^t\|\Delta u\|_{L^2}^2 d\tau\\ &\quad\le C\int_0^t \|\partial_2 u_2\|_{L^{\alpha_3}}^{\frac{2}{r-2}}\|\nabla u\|_{L^2}^2 d\tau+C\int_0^t\|u_3\|_{L^{\alpha_1}}^{\frac{2\alpha_1}{2\alpha_1-3}}\|\nabla u\|_{L^2}^2 d\tau+\|\nabla u(0)\|_{L^2}^2.\end{aligned} \tag{2.65}$$

We claim $\frac{2}{r-2}=\beta_3$. In fact, from (2.50), we have

$$\gamma=\frac{-3+\sqrt{24\alpha_3^2-24\alpha_3+9}}{2(\alpha_3-1)}$$

and then by the definition (2.62), one has

$$\frac{2}{\frac{2}{r-2}}+1=r-1=\gamma-\frac{\gamma}{\alpha_3}=\frac{-3+\sqrt{24\alpha_3^2-24\alpha_3+9}}{2\alpha_3}, \tag{2.66}$$

Comparing (2.66) with (2.56), we prove the claim. Therefore, we can apply Gronwall's inequality to (2.65), and by condition (1.11), (1.14) and (1.15) to get that the H^1 norm of the strong solution u is bounded on the maximal interval of existence $(0,T^*)$. The proof of this Theorem is completed.

Remark 2.2. In the process of the proof, if we want to prove the case of $i=3$ when $j=2$. For the first part, the inequality (2.42) may be replaced by

$$K_2(t)\le C\|u_2\|_{L^2}^{\frac{r-1}{r}}\|\partial_3 u_2\|_{L^{\frac{2}{3-r}}}^{\frac{1}{r}}\|\nabla u\|_{L^2}^{\frac{r-2}{r}}\|\Delta u\|_{L^2}^{\frac{r+2}{r}}, \tag{2.67}$$

and (2.43) becomes

$$K_2(t)\le C\|u_2\|_{L^2}^{\frac{4(\alpha_2-1)}{\alpha_2-2}}\|\partial_3 u_2\|_{L^{\alpha_2}}^{\frac{2\alpha_2}{\alpha_2-2}}\|\nabla u\|_{L^2}^2+\frac{\nu}{4}\|\Delta u\|_{L^2}^2. \tag{2.68}$$

If we want to prove the case of $j = 1$, we shall give an alternative proof of the term I_2, also by the incompressibility condition, one has

$$\begin{aligned} I_2 &= -\int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_1 \partial_3 u_1 + \partial_3 u_2 \partial_2 u_1 \partial_3 u_1 + \partial_3 u_3 \partial_3 u_1 \partial_3 u_1 dx \\ &\quad - \int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_2 \partial_3 u_2 + \partial_3 u_2 \partial_2 u_2 \partial_3 u_2 + \partial_3 u_3 \partial_3 u_2 \partial_3 u_2 dx \\ &= -\int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_1 \partial_3 u_1 + \partial_3 u_2 \partial_2 u_1 \partial_3 u_1 + \partial_3 u_3 \partial_3 u_1 \partial_3 u_1 dx \\ &\quad - \int_{\mathbb{R}^3} \partial_3 u_1 \partial_1 u_2 \partial_3 u_2 + \partial_3 u_2 \left(-\partial_1 u_1 - \partial_3 u_3\right) \partial_3 u_2 + \partial_3 u_3 \partial_3 u_2 \partial_3 u_2 dx \\ &\leq \int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx + \int_{\mathbb{R}^3} |u_1||\nabla u||\nabla^2 u| dx. \end{aligned}$$

and then we obtain

$$\begin{aligned} &\frac{1}{2}\frac{d}{dt}\|\nabla u\|_{L^2}^2 + \nu\|\Delta u\|_{L^2}^2 \\ &\qquad \leq C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx + C\int_{\mathbb{R}^3} |u_1||\nabla u||\nabla^2 u| dx, \\ &\qquad = K_1(t) + K_2(t), \end{aligned} \tag{2.69}$$

by which one can prove the case of $i \neq j = 1$ and $i = j = 1$. The term $K_1(t)$ is same to (2.44). As for the second term $K_2(t)$, we shall give the inequality (2.42) as the following for $i \neq j = 1$,

$$K_2(t) \leq C\|u_1\|_{L^2}^{\frac{r-1}{r}} \|\partial_i u_1\|_{L^{\frac{2}{3-r}}}^{\frac{1}{r}} \|\nabla u\|_{L^2}^{\frac{r-2}{r}} \|\Delta u\|_{L^2}^{\frac{r+2}{r}}, \ i = 2 \text{ or } 3, \tag{2.70}$$

and then get the corresponding form of (2.43). As for $i = j = 1$, we will use $|u_1|^{\gamma-2}u_1$, as test function in the equation for u_1, and we can get the similar results to (2.57). By the same process to prove the case of $i = j = 1$.

Proof of Theorem 1.7 Give the same process as in the Theorem 1.3, we have

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|\nabla u\|_2^2 + \nu\|\Delta u\|_2^2 &\leq C\int_{\mathbb{R}^3} |\partial_3 u_2||\nabla u|^2 dx + C\int_{\mathbb{R}^3} |u_3||\nabla u||\nabla^2 u| dx \\ &\quad + C\int_{\mathbb{R}^3} |\partial_3 u_2||\Delta u||u| dx + C\int_{\mathbb{R}^3} |\partial_3 u_3||\Delta u||u| dx \\ &= L_1 + L_2 + L_3 + L_4. \end{aligned} \tag{2.71}$$

Here, we only prove the case of $\partial_3 u_2, \partial_3 u_3$. For the case of $\partial_3 u_1, \partial_3 u_3$, we will begin with (2.39), and form which we can give the similar proof.

We estimate L_i, $i = 1, 2, 3, 4$ one by one. Firstly, for L_1 we have

$$L_1 \leq C\|\partial_3 u_2\|_{L^{\alpha_1}} \|\nabla u\|_{L^{r_1}}^2, \tag{2.72}$$

where r_1 satisfies

$$\frac{1}{\alpha_1} + \frac{2}{r_1} = 1 \Rightarrow r_1 = \frac{2\alpha_1}{\alpha_1 - 1}, \tag{2.73}$$

since $2 \leq \alpha_1 \leq 3$, we have $3 \leq r_1 \leq 4$, by Gagliardo-Nirenberg and Young's inequalities, on has

$$\begin{aligned} L_1 &\leq C\|\partial_3 u_2\|_{L^{\alpha_1}} \|\nabla u\|_{L^2}^{\frac{6-r_1}{r_1}} \|\Delta u\|_{L^2}^{\frac{3r_1-6}{r_1}} \\ &\leq C\|\partial_3 u_2\|_{L^{\alpha_1}}^{\frac{2r_1}{6-r_1}} \|\nabla u\|_{L^2}^2 + \frac{\nu}{4}\|\Delta u\|_{L^2}^2 \\ &= C\|\partial_3 u_2\|_{L^{\alpha_1}}^{\frac{2\alpha_1}{2\alpha_1-3}} \|\nabla u\|_{L^2}^2 + \frac{\nu}{4}\|\Delta u\|_{L^2}^2. \end{aligned} \tag{2.74}$$

As for L_3, let r_2 satisfy

$$\frac{1}{\alpha_1} + \frac{1}{r_2} = \frac{1}{2}, \tag{2.75}$$

then we have

$$L_3 \leq C\|\partial_3 u_2\|_{L^{\alpha_1}} \|\Delta u\|_{L^2} \|u\|_{L^{r_2}}. \tag{2.76}$$

From the fact that $2 \leq \alpha_1 \leq 3$, we have $r_2 \geq 6$. By the Gagliardo-Nirenberg inequality,

$$\|v\|_{L^r} \leq \|\nabla v\|_{L^{\frac{3r}{3+r}}}, \quad \frac{3}{2} \leq r < \infty. \tag{2.77}$$

Therefore, we have

$$\|u\|_{L^{r_2}} \leq \|\nabla u\|_{L^{r_3}}, \tag{2.78}$$

where $r_3 = 3r_2/(3+r_2)$ with $2 \leq r_3 \leq 3$. Thus, we have

$$\frac{1}{\alpha_1} + \frac{1}{r_3} = \frac{5}{6} \Rightarrow r_3 = \frac{6\alpha_1}{5\alpha_1 - 6}, \tag{2.79}$$

and applying Young's inequality, (2.76) becomes

$$\begin{aligned} L_3 &\leq C\|\partial_3 u_2\|_{L^{\alpha_1}} \|\Delta u\|_{L^2} \|\nabla u\|_{L^{r_3}}. \\ &\leq C\|\partial_3 u_2\|_{L^{\alpha_1}} \|\nabla u\|_{L^2}^{\frac{6-r_3}{2r_3}} \|\Delta u\|_{L^2}^{\frac{5r_3-6}{2r_3}}. \\ &\leq C\|\partial_3 u_2\|_{L^{\alpha_1}}^{\frac{4r_3}{6-r_3}} \|\nabla u\|_{L^2}^2 + \frac{\nu}{8}\|\Delta u\|_{L^2}^2. \\ &= C\|\partial_3 u_2\|_{L^{\alpha_1}}^{\frac{2\alpha_1}{2\alpha_1-3}} \|\nabla u\|_{L^2}^2 + \frac{\nu}{8}\|\Delta u\|_{L^2}^2. \end{aligned} \tag{2.80}$$

The term L_4 is estimated in the same way and we get

$$L_4 \leq C\|\partial_3 u_3\|_{L^{\alpha_2}}^{\frac{2\alpha_2}{2\alpha_2-3}} \|\nabla u\|_{L^2}^2 + \frac{\nu}{8}\|\Delta u\|_{L^2}^2. \tag{2.81}$$

Therefore, integrating on time and absorbing the last term in (2.74), (2.80) and (2.81), we get

$$\begin{aligned} &\|\nabla u\|_2^2 + \frac{3\nu}{2}\int_0^t \|\Delta u\|_2^2 d\tau \\ &\quad\leq \|\nabla u(0)\|_2^2 + C\int_0^t \|\partial_3 u_2\|_{\alpha_1}^{\frac{2\alpha_1}{2\alpha_1-3}} \|\nabla u\|_2^2 d\tau \\ &\qquad + C\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\frac{2\alpha_2}{2\alpha_2-3}} \|\nabla u\|_2^2 d\tau + C\int_0^t L_2(\tau) d\tau. \end{aligned} \tag{2.82}$$

As for the estimation of L_2, we give the same proof as the case of $i=j$ in Theorem 1.4, in which u_2 is replaced by u_3, $\partial_2 u_2$ is replaced by $\partial_3 u_3$ and α_3, β_3 is replaced by α_2, β_2. Finally, we get

$$\begin{aligned}\int_0^t L_2(\tau)d\tau &\le C\int_0^t \|u_3\|_\gamma^{\frac{2(r-1)}{r-2}}\|\partial_3 u_3\|_{\alpha_2}^{\frac{2}{r-2}}\|\nabla u\|_2^2 d\tau + \frac{\nu}{2}\int_0^t \|\Delta u\|_2^2 d\tau \\ &\le C\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\frac{2}{r-2}}\|\nabla u\|_2^2 d\tau + \frac{\nu}{2}\int_0^t \|\Delta u\|_2^2 d\tau,\end{aligned} \tag{2.83}$$

where $\alpha_2 \ge 2$ and

$$\begin{cases} \dfrac{1}{\gamma} = \dfrac{3+\sqrt{24\alpha_2^2-24\alpha_2+9}}{12\alpha_2}, \\ \mu = \dfrac{\alpha_2\gamma}{2\alpha_2-\gamma}, \\ r = \dfrac{(\gamma+1)\alpha_2-\gamma}{\alpha_2}. \end{cases} \tag{2.84}$$

Inserting (2.83) into (2.82) and absorbing the last term in (2.83), note the boundedness of $\|u_3\|_\gamma$, we have

$$\begin{aligned}&\|\nabla u\|_2^2 + \nu\int_0^t \|\Delta u\|_2^2 d\tau \\ &\quad\le \|\nabla u(0)\|_2^2 + C\int_0^t \|\partial_3 u_2\|_{\alpha_1}^{\frac{2\alpha_1}{2\alpha_1-3}}\|\nabla u\|_2^2 d\tau \\ &\qquad + C\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\frac{2\alpha_2}{2\alpha_2-3}}\|\nabla u\|_2^2 d\tau + C\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\frac{2}{r-2}}\|\nabla u\|_2^2 d\tau.\end{aligned} \tag{2.85}$$

We also can check that $\frac{2}{r-2} = \beta_2$ (see (2.66)), where $\beta_2 = \frac{2\mu}{3-\mu}$. Denote that

$$\frac{1}{\beta'} = \frac{2\alpha_2-3}{2\alpha_2}.$$

Also from (2.66)

$$r = \frac{-3+\sqrt{24\alpha_2^2-24\alpha_2+9}}{2\alpha_2} + 1.$$

We see that

$$\begin{aligned}\frac{1}{\beta'} - \frac{1}{\beta_2} &= \frac{2\alpha_2-3}{2\alpha_2} - \frac{r-2}{2} \\ &= \frac{(4-r)\alpha_2-3}{2\alpha_2} \\ &= \frac{\frac{6\alpha_2+3-\sqrt{24\alpha_2^2-24\alpha_2+9}}{2}-3}{2\alpha_2} \\ &= \frac{6\alpha_2-3-\sqrt{24\alpha_2^2-24\alpha_2+9}}{4\alpha_2} \\ &= \frac{12\alpha_2(\alpha_2-1)}{4\alpha_2\left((6\alpha_2-3)+\sqrt{24\alpha_2^2-24\alpha_2+9}\right)} > 0, \ \forall\ \alpha_2 \ge 2.\end{aligned} \tag{2.86}$$

From above inequality, we know $\beta' < \beta_2$ and hance $\beta'/\beta_2 < 1$. Therefore, applying Hölder's inequality, and by energy inequality, one has

$$\begin{aligned}
\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\frac{2\alpha_2}{2\alpha_2-3}} \|\nabla u\|_2^2 d\tau &= \int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\beta'} \|\nabla u\|_2^2 d\tau \\
&= \int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\beta'} \|\nabla u\|_2^{\frac{2\beta'}{\beta_2}} d\tau \|\nabla u\|_2^{\frac{2(\beta_2-\beta')}{\beta_2}} d\tau \\
&\le C\left(\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\beta_2} \|\nabla u\|_2^2 d\tau\right)^{\frac{\beta'}{\beta_2}} \left(\int_0^t \|\nabla u\|_2^2 d\tau\right)^{\frac{\beta_2-\beta'}{\beta_2}} \qquad (2.87) \\
&\le C\left(\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\beta_2} \|\nabla u\|_2^2 d\tau\right)^{\frac{\beta'}{\beta_2}} \\
&\le C\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\frac{2}{r-2}} \|\nabla u\|_2^2 d\tau + C.
\end{aligned}$$

Finally, we get

$$\begin{aligned}
\|\nabla u\|_2^2 \quad &+\nu \int_0^t \|\Delta u\|_2^2 d\tau \\
&\le \|\nabla u(0)\|_2^2 + C\int_0^t \|\partial_3 u_2\|_{\alpha_1}^{\frac{2\alpha_1}{2\alpha_1-3}} \|\nabla u\|_2^2 d\tau \qquad (2.88) \\
&\quad +C\int_0^t \|\partial_3 u_3\|_{\alpha_2}^{\frac{2}{r-2}} \|\nabla u\|_2^2 d\tau + C.
\end{aligned}$$

By using Gronwall's inequality to (2.88), and by condition (1.18), (1.19) and (1.20), we get that the H^1 norm of the strong solution u is bounded on the maximal interval of existence $(0, T^*)$. The proof of the case of $\partial_3 u_2, \partial_3 u_3$ is completed.

Proof of Theorem 1.10 Without loss of generality, we assume $j = 3, k = 3$. For every

$$\alpha \in (\frac{9}{5}, \infty), \tag{2.89}$$

we set

$$\begin{cases}
\dfrac{1}{\mu} = \dfrac{1 - \frac{12}{\alpha} + \sqrt{\frac{144}{\alpha^2} - \frac{264}{\alpha} + 289}}{24}, \\
q_2 = \dfrac{2\alpha\mu}{\alpha+\mu}, \\
r = \dfrac{2\mu\alpha - \mu + \alpha}{\alpha+\mu}.
\end{cases} \tag{2.90}$$

From (2.90), we see that μ is a decreasing function of α, and by (2.89), we have

$$\frac{4}{3} < \mu < 3, \tag{2.91}$$

and (2.90) follows

$$\frac{1}{\mu} + \frac{1}{\alpha} + \frac{q_2-2}{q_2} = 1. \tag{2.92}$$

On the other hand, also from (2.90), we note that μ satisfies

$$\alpha = \frac{12\mu + 5\mu^2}{6\mu^2 + \mu - 12}, \tag{2.93}$$

and

$$\frac{12\mu + 5\mu^2}{6\mu^2 + \mu - 12} - \frac{\mu}{\mu - 1} = \frac{\mu^2(6-\mu)}{(6\mu^2 + \mu - 12)(\mu - 1)} > 0 \text{ with } \frac{4}{3} < \mu < 3.$$

Combining (2.93) and above inequality, we have $\alpha > \frac{\mu}{\mu-1}$, and hance (2.90) implies $q_2 > 2$. We choose

$$\beta = \frac{2\mu}{3-\mu}, \tag{2.94}$$

then from (2.90), (2.92) and (2.94), we can compute that

$$\frac{2}{\beta} + 1 = \frac{3}{\mu} = \frac{\alpha + \sqrt{289\alpha^2 - 264\alpha + 144}}{8\alpha} - \frac{3}{2\alpha} \Rightarrow \frac{3}{2\alpha} + \frac{2}{\beta} = g(\alpha). \tag{2.95}$$

We denote

$$V_1(t) = \int_0^t \|\partial_3 u_3\|_{L^\alpha}^{\beta} \|\nabla u\|_{L^2}^2 d\tau = \int_0^t \|\partial_3 u_3\|_{L^\alpha}^{\frac{2\mu}{3-\mu}} \|\nabla u\|_{L^2}^2 d\tau. \tag{2.96}$$

Next, we give a estimates on u_3. We use $|u_3|^{q_2-2}u_3$ as test function in the equation (1.1) for u_3. Similar to (2.57), we have

$$\|u_3\|_{L^{q_2}}^2 \le \|u_3(0)\|_{L^{q_2}}^2 + C + C\int_0^t \|\partial_3 u_3\|_{L^\alpha}^{\frac{2\mu}{3-\mu}} d\tau. \tag{2.97}$$

By the condition (1.24), (2.90) and (2.91), we have $q_2 < 6$. Note that $\|u_3(0)\|_{q_2} < C$ for some $C > 0$, we get

$$u_3 \in L^\infty(0, T; L^{q_2}(\mathbb{R}^3)). \tag{2.98}$$

By (2.90), we have $r = \frac{(q_2+1)\alpha - q_2}{\alpha}$, and $r > 2$, for such q_2, r, α, we can apply Lemma 2.3 in [9], and we have

$$\begin{aligned} &\|\nabla u\|_{L^2}^2 + \frac{\nu}{2}\int_0^t \|\Delta u\|_{L^2}^2 d\tau \\ &\le \|\nabla u(0)\|_{L^2}^2 + C\int_0^t \|u_3\|_{L^{q_2}}^{\frac{2(r-1)}{r-2}} \|\partial_3 u_3\|_{L^\alpha}^{\frac{2}{r-2}} \|\nabla u\|_{L^2}^2 d\tau \\ &+C\int_0^t \|u_3\|_{L^{q_2}}^{\frac{8(r-1)}{3(r-2)}} \|\partial_3 u_3\|_{L^\alpha}^{\frac{8}{3(r-2)}} \|\nabla u\|_{L^2}^2 d\tau + C. \end{aligned} \tag{2.99}$$

Moreover, r satisfies

$$\frac{8}{3(r-2)} = \frac{8(\mu+\alpha)}{3(2\mu\alpha - 3\mu - \alpha)} > 0.$$

From (2.93), we have (note that $2\mu\alpha - 3\mu - \alpha > 0$)

$$\frac{8(\mu+\alpha)}{3(2\mu\alpha - 3\mu - \alpha)} - \frac{2\mu}{3-\mu} = \frac{2[12\mu + 5\mu^2 + (-6\mu^2 - \mu + 12)\alpha]}{3(2\mu\alpha - 3\mu - \alpha)(3-\mu)} = 0.$$

Combining (2.98) and (2.99), and the fact $\frac{2}{r-2} < \frac{8}{3(r-2)}$, we have

$$\|\nabla u\|_{L^2}^2 + \nu\int_0^t \|\Delta u\|_{L^2}^2 d\tau \le CV_1(t) + \|\nabla u(0)\|_{L^2}^2 + C,$$

and end the proof for $\alpha \in \left(\frac{9}{5}, \infty\right)$ by using Gronwall's inequality.

Acknowledgement

The second author would like to thank Dr. Ting Zhang for his helpful suggestions. This work is supported partially by NSFC 11271322, 10931007, and Zhejiang NSF of China Z6100217.

References

[1] H.O. Bae, H. J. Choe, A Regularity Criterion for the Navier-Stokes Equations. Communications in Partial Differential Equations, 32: 1173-1187 (2007).

[2] H. Beirão da Veiga, Concerning the regularity problem for the solutions of the Navier-Stokes equations, C.R. Acad. Sci. Paris, t.321. Série I (1995), 405-408.

[3] L.C. Berselli, On a regularity criterion for the solutions to the 3D Navier-Stokes equations, Dierential Integral Equations 15, 1129-1137 (2002).

[4] L. Caffarelli, R. Kohn, L. Nirenberg, Partial regularity of suitable weak solutions of the NavierCStokes equations Commun. Pure Appl. Math. 35 771-831 (1982).

[5] C.S. Cao, E.S. Titi, Global Regularity Criterion for the 3D Navier-Stokes Equations Involving One Entry of the Velocity Gradient Tensor. Arch. Rational Mech.Anal. 202(2011) 919-932.

[6] C.S. Cao, E.S. Titi, Regularity criteria for the three-dimensional Navier-Stokes equations, Indiana Univ. Math. J. 57(6), 2643-2661 (2008).

[7] D. Chae, H. J. Choe: Regularity of solutions to the Navier-Stokes equation. Electron. J. Differential Equations 5 (1999), 1-7.

[8] P. Constantin, C. Foias, Navier-Stokes Equations. The University of Chicago Press, Chicago, 1988.

[9] D.Y. Fang, C.Y. Qian, The regularity criterion for 3D Navier-Stokes Equations involving one velocity gradient component. Nonlinear analysis 78 (2013) 86-103.

[10] G.P. Galdi, An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Vol. I, II. Springer, New York, 1994.

[11] E. Hopf, Über die anfang swetaufgabe für die hydrodynamischer grundgleichungan, Math. Nach. 4 (1951), 213-231.

[12] I. Kukavica, M. Ziane, One component regularity for the Navier-Stokes equations, Nonlinearity 19, 453-469 (2006).

[13] I. Kukavica, M. Ziane, Navier-Stokes equations with regularity in one direction. J. Math. Phys. 48(6), 065203, 10 pp (2007)

[14] O.A. Ladyzhenskaya, The Boundary Value Problems of Mathematical Physics. Springer, Berlin, 1985.

[15] J. Leray, Sur le mouvement d'um liquide visqieux emlissant l'space, Acta Math. 63 (1934), 193-248.

[16] J. Neustupa, A. Novotny, and P. Penel, An interior regularity of a weak solution to the Navier-Stokes equations dependence on one component of velocity, Topics in Mathematical Fluid Mechanics, Quaderni di Matematica, Dep Math. Seconda University, Napoli, Caserta, Vol. 10, pp. 163-183, 2002; see also: A remark to interior regularity of suitable weak solution to the Navier-Stokes equations, CIM Preprint No. 25, 1999.

[17] P. Penel, M. Pokorný, On anisotropic regularity criteria for the Solutions to 3D Navier-Stokes equations, J. Math. Fluid Mech. 13 (2011), 341-353.

[18] P.Penel, M. Pokorný, Some new regularity criteria for the Navier-Stokes equations containing gradient of the velocity, Appl.Math. 49, 483-493 (2004).

[19] M. Pokorný, On the result of He concerning the smoothness of solutions to the Navier-Stokes equations. Electron. J. Differ. Equ. 11, 1-8 (2003).

[20] G. Prodi, Un teorema di unicità per le equazioni di Navier-Stokes, Ann. Mat. Pura Appl. IV 48,173 (1959).

[21] J. Serrin, The initial value problems for the Navier-Stokes equations, in Nonlinear Problems, edited by Langer, R. E. (University of Wisconsin Press, Madison, WI, 1963).

[22] H. Sohr, The Navier-Stokes Equations, An Elementary Functional Analytic Approach. Birkhäuser Verlag, Basel, 2001.

[23] E. M. Stein, Singular integrals and diffierentiability properties of functions. Princeton Mathematical Series, No. 30 Princeton University Press, Princeton, NJ 1970.

[24] R. Temam, Navier-Stokes Equations, Theory and Numerical Analysis. North-Holland, 1984.

[25] Y. Zhou, A new regularity criterion for weak solutions to the Navier-Stokes equations, J. Math. Pures Appl. 84, 1496-1514 (2005).

[26] Y. Zhou, M. Pokorný, On the regularity of the solutions of the Navier-Stokes equations via one velocity component. Nonlinearity 23, 1097-1107 (2010).

[27] Y. Zhou, M. Pokorný, On a regularity criterion for the Navier-Stokes equations involving gradient of one velocity component. J. Math Phys. 50, 123514 (2009). doi:10.1063/1.3268589.